# **New specimens from Mid-Cretaceous Myanmar** amber illuminate the phylogenetic placement of Lagonomegopidae (Arachnida: Araneae)

XIANGBO GUO<sup>1</sup>, PAUL A. SELDEN FLS<sup>1,2,3,\*</sup> and DONG REN<sup>1,\*,0</sup>

<sup>1</sup>College of Life Sciences and Academy for Multidisciplinary Studies, Capital Normal University, 105 Xisanhuanbeilu, Haidian District, Beijing, 100048, China <sup>2</sup>Department of Geology, University of Kansas, 1457 Jayhawk Boulevard, Lawrence KS 66045, USA <sup>3</sup>Natural History Museum, Cromwell Road, London SW7 5BD, UK

Received 16 January 2021; revised 8 March 2021; accepted for publication 9 April 2021

New lagonomegopid spiders are described from Mid-Cretaceous Myanmar (Burmese) amber. Two new genera and species based on single specimens, Scopomegops fax gen. & sp. nov. and Hiatomegops spinalis gen. & sp. **nov.** are described. Two specimens belonging to *Lineaburmops beigeli* are further described. Additionally, after re-examining the holotype of Odontomegops titan, a detailed description of its basal ventral abdomen is added here. A phylogenetic analysis was performed to investigate the phylogenetic placement of Lagonomegopidae. A matrix of 79 morphological characters, scored for six lagonomegopid taxa and 26 non-lagonomegopid taxa, was analysed through parsimony and Bayesian phylogenetic inference. Our results recover extant Palpimanoidea as a monophyletic group and partly suggest that Lagonomegopidae is the sister-group to extant Palpimanoidea. The external sexual organs, retrolateral tibial apophysis on the male palp and tracheal spiracle in lagonomegopids are discussed.

ADDITIONAL KEYWORDS: fossil - Mesozoic - Palpimanoidea - retrolateral tibial apophysis - tracheal spiracle.

# INTRODUCTION

Spiders (Araneae) are one of the most diverse animal groups and an important component of every terrestrial ecosystem, with more than 49 000 described extant species and 1300 described fossil species (Dunlop et al., 2020; World Spider Catalog, 2021). Many recent phylogenetic studies, which improved our knowledge of the systematics and evolution of spiders, were based on the morphological, behavioural and molecular data of extant taxa (e.g. Griswold et al., 2005; Bond et al., 2014; Fernández et al., 2014; Ramírez, 2014; Polotow et al., 2015; Garrison et al., 2016; Dimitrov et al., 2017; Wheeler et al., 2017; Azevedo et al., 2018; Wood et al., 2018; Opatova et al., 2020). Extinct spider taxa have usually been used for anchoring the age of particular nodes in the phylogenetic trees depicting

relationships among extant taxa (e.g. Dimitrov et al., 2013; Benavides et al., 2017; Fernández et al., 2018; Li & Li, 2018; Shao & Li, 2018; Magalhaes et al., 2020). By contrast, only several studies used fossil spiders as terminal taxa in the phylogenetic analysis (Wood et al., 2012, 2013, 2015; Wood, 2017; Selden et al., 2020). Therefore, the phylogenetic placement of spider fossils is often ambiguous (Magalhaes et al., 2020).

The fossil spider family Lagonomegopidae, widespread in the Cretaceous period (Park et al., 2019), was deemed not to be a stem group of some extant lineage because of its highly derived morphology, and thus it was likely to represent an extinct lineage (Magalhaes *et al.*, 2020). It was incorporated into the superfamily Palpimanoidea primarily based on the presence of peg teeth and the absence of true teeth on the cheliceral promargin, as well as the trichobothrial pattern and spineless legs (Eskov & Wunderlich, 1995). However, as more and more lagonomegopids were reported, some characters weakening the support for its superfamilial placement in Palpimanoidea, such as tarsal trichobothria and feathery setae, were

<sup>\*</sup>Corresponding authors. E-mail: rendong@mail.cnu.edu.cn; selden@ku.edu

<sup>[</sup>Version of record, published online 16 June 2021; http://zoobank.org/urn:lsid:zoobank. org:pub:8FEFC59E-7194-43DD-A633-6359A8936921]

described (Penney, 2004, 2005; Guo *et al.*, 2020). The phylogenetic placement of Lagonomegopidae was thus considered controversial (Pérez-de la Fuente *et al.*, 2013; Guo *et al.*, 2020).

Forster & Platnick (1984) based an expanded superfamily Palpimanoidea, including ten families, on two cheliceral features: the presence of peg teeth on the promargin and an elevated cheliceral gland mound. But some subsequent studies did not support this expanded Palpimanoidea (Schütt, 2000; Griswold et al., 2005; Rix et al., 2008; Dimitrov et al., 2012). Wood et al. (2012) performed a phylogenetic analysis using both morphological and molecular data. This total-evidence analysis resulted in a delimitation of the superfamily Palpimanoidea to contain five families: Archaeidae Koch & Berendt, 1854, Palpimanidae Thorell, 1870, Stenochilidae Thorell, 1873, Huttoniidae Simon, 1893 and Mecysmaucheniidae Simon, 1895. Although another study based on six genetic markers suggested that Palpimanoidea is paraphyletic (Wheeler et al., 2017), two recent studies using transcriptomic data and DNA sequences from Ultra-Conserved Elements and exonic loci recovered a monophyletic Palpimanoidea (Fernández et al., 2018; Wood et al., 2018). All these recent analyses (Wood et al., 2012, 2013, 2018; Wheeler et al., 2017; Fernández et al., 2018) considered Palpimanoidea as a possible sister-group of Entelegynae, a more diverse and derived spider group. Thus, palpimanoids are an important group for understanding spider evolution (Wood et al., 2018).

Except Lagonomegopidae, there are three fossil families (Spatiatoridae Petrunkevitch, 1942, Micropalpimanidae Wunderlich, 2008 and Vetiatoridae Wunderlich, 2017) and four fossil genera (family uncertain; Sinaranea Selden et al., 2008, Seppo Selden & Dunlop, 2014, Caestaranea Selden et al., 2020 and Onychopalpus Selden et al., 2020) placed in the superfamily Palpimanoidea. But only Sinaranea, Caestaranea and Onychopalpus have been analysed cladistically for their phylogenetic positions (Selden et al., 2020). Fossils can have crucial implications for understanding spider evolution; indeed, detailed examination of spider fossils using modern methods is critical to better understand and interpret the phenotypic features and phylogenetic affinities of extinct taxa (Dimitrov & Hormiga, 2021). Moreover, exploring the behaviours and lifestyles of fossil taxa can provide a window into the sophisticated ecosystems in geological history (Lin et al., 2019; Zhao et al., 2020; Gao et al., 2021), but such research focusing on fossil spiders is relatively scarce.

Herein, two new genera and species, as well as new specimens of *Lineaburmops beigeli* Wunderlich, 2015, are described from Burmese amber. A supplementary description of *Odontomegops titan* Guo & Selden, 2020 is provided after re-examining its holotype. Phylogenetic analyses based on morphological data were carried out to test the phylogenetic placement of Lagonomegopidae. We recovered extant Palpimanoidea as a monophyletic group and our implied weighted analyses suggests that Lagonomegopidae is the sister-group to extant Palpimanoidea.

## MATERIAL AND METHODS

#### MATERIAL AND TERMINOLOGY

The amber specimens investigated in this paper are from Tanai Village in the Hukawng Valley, Myitkyina District of Kachin State, Myanmar (Cruickshank & Ko, 2003: fig. 1). The amber-bearing deposits have been dated to the earliest Cenomanian, c. 98.8  $\pm$  0.6 Mya, based on U– Pb radiometric dating of zircons from the volcaniclastic matrix (Shi *et al.*, 2012). All specimens are housed at the fossil collection of the Key Lab of Insect Evolution & Environmental Changes, at the College of Life Sciences, Capital Normal University (CNUB; Dong Ren, curator), in Beijing, China. They were acquired by Fangyuan Xia before 2013 and donated for this study in 2015.

Preparation and imaging methods follow Selden & Penney (2017). The photographs were taken with a Nikon SMZ 25 and an attached Nikon DS-Ri 2 digital camera system, as well as a Nikon ECLIPSE Ni and an attached Nikon DS-Ri 2 digital camera system. Micro-CT scanning of specimen CNU-ARA-MA2020001 was carried out with a Micro-CT (HeliScan micro-CT, Thermos Fisher Scientific), located at the Department of Geology at the University of Kansas. The voltage of the Micro-CT scanner was 100 KV. The three-dimensional structure of the male palp of CNU-ARA-MA2020001 was reconstructed using 3D Slicer 4.10.2. The line drawings were prepared with Adobe Illustrator CS6 and Adobe Photoshop CC, the images were processed by Adobe Photoshop CC. Measuring method follows Selden et al. (2016); measurements of coxae and trochanters are imprecise and uninformative. All measurements are in mm.

Leg formula indicates the length of each leg relative to the others, longest to shortest. Abbreviations: I, II, III, IV, leg numbers; ALE, anterior lateral eye(s); ALS, anterior lateral spinneret(s); AME, anterior median eye(s); at, anal tubercle; C, conductor; cx, coxa; ef, epigastric furrow; E, embolus; fe, femur; MA, median apophysis; mt, metatarsus; P, protrusion(s); pa, patella; PLE, posterior lateral eye(s); PLS, posterior lateral spinneret(s); PME, posterior median eye(s); PMS, posterior median spinneret(s); RTA, retrolateral tibial apophysis; ST, subtegulum; ta, tarsus; ti, tibia; tr, trochanter; ts, tracheal spiracle.

#### TAXON SAMPLING AND CHARACTER CHOICE

The discovery of new lagonomegopid fossils with wellpreserved morphological details (Park *et al.*, 2019; Guo *et al.*, 2020) provided incentive for our analysis of the phylogenetic placement of Lagonomegopidae. We sampled six terminal taxa from Burmese amber (five taxa) and the Jinju Formation of Korea (one taxon) representing Lagonomegopidae. These taxa comprise adult males offour species: *Lineaburmops beigeli*, *Jinjumegops dalingwateri* Park *et al.*, 2019, *Odontomegops titan* and *Scopomegops fax*, one adult female: *Hiatomegops spinalis*, and one subadult male: Lagonomegopidae *indet*. described by Guo *et al.* (2020). They were selected because the specimens are well preserved and show some characters which were reported for the first time in Lagonomegopidae.

With reference to previous phylogenetic analyses of Araneomorphae (Griswold *et al.*, 2005; Wood *et al.*, 2012), we sampled an additional 26 non-lagonomegopid taxa representing the major Araneae clades, with the most basal spider family Liphistiidae and mygalomorph family Antrodiaetidae as the outgroups. Five living families of Palpimanoidea (Wood *et al.*, 2012; Wheeler *et al.*, 2017; Fernández *et al.*, 2018), as well as *Mimetus* Hentz, 1832 and *Pararchaea* Forster, 1955, which have peg teeth on the cheliceral promargin but belong to the superfamily Araneoidea (Wood *et al.*, 2012; Garrison *et al.*, 2016; Wheeler *et al.*, 2017; Fernández *et al.*, 2018), were included in this study.

Sixty-seven characters were extracted from previous analyses (Griswold et al., 2005; Wood et al., 2012) and 12 new characters were added for this analysis. Character states of non-lagonomegopid spiders were scored using data from other studies (Forster & Platnick, 1984; Griswold, 1990; Wesołowska, 1999; Griswold et al., 2005; Hendrixson & Bond, 2007; Wood et al., 2012; Ramírez, 2014; Murphy & Roberts, 2015; Xu et al., 2015), as well as from SEM images acquired from other arachnology labs as part of the NSF Assembling the Tree of Life - Spiders project (http://www.morphbank.net). MESQUITE v.3.6 (Maddison & Maddison, 2019) was used to assemble the character matrix. These characters were coded as unordered. Inapplicable and unknown characters were respectively coded as '-' and '?'. Descriptions of character states are given in the Supporting Information (File S1). All the taxa used can be found in the Supporting Information (File S2). The data matrix is given in the Supporting Information (Files S2, S3).

#### PHYLOGENETIC ANALYSIS

Phylogenetic analyses were carried out using maximum parsimony and Bayesian methods. The optimal trees under the parsimony criterion were sought in TNT v.1.5 (Goloboff *et al.*, 2008a; Goloboff & Catalano, 2016) using the traditional search option under the following parameters: tree bisection reconnection (TBR) branchswapping algorithm with 1000 replications saving 100 trees per replicate. Searches were performed under equal weights and implied weights (Goloboff, 1993; Goloboff et al., 2008b). In our implied weights analysis, we used the methodology proposed by Mirande (2009) to set the value of concavity constant K, which determines how strongly the analysis will weigh against homoplasy. The TNT script developed by Mirande (2009: appendices S3, S4) was used (script commands aaa 2 16 50 95 7) to find the most fit trees under a wide range of *K* values. The best trees under different values of K were compared using SPR distances, the most stable tree had the highest average SPR similarity, which defined the best K value (Goloboff, 2008). Clade support was estimated within TNT through symmetric resampling (Goloboff et al., 2003; absolute frequencies, change probability 33%, 1000 replicates) and by Bremer (1994) support (using a script that comes with the TNT package). The characters were mapped in WinClada (Nixon, 2002).

Bayesian analyses were performed in MrBayes v.3.2.7 (Ronquist & Huelsenbeck, 2003; Ronquist *et al.*, 2012) using a NEXUS file (including MrBayes' commands) provided by Selden *et al.* (2020: supplementary data). The analysis employed the Lewis (2001) discrete (morphology) model. Two runs of four chains, three of which were heated, were run for a total of 5 000 000 generations, with sampling every 500 generations and the first 25% discarded as burn-in. A majority-rule consensus tree was computed with posterior probabilities for each node.

#### RESULTS

#### PHYLOGENETIC RESULTS

The equal weights analysis found 50 most parsimonious trees with 214 steps [consistency index (CI) = 41, retention index (RI) = 62]. The strict consensus of these trees (Supporting Information, Fig. S1) was unresolved regarding the relationships within Araneomorphae. The Bayesian analysis had a similar result (Supporting Information, Fig. S2). Extant Palpimanoidea were recovered as a monophyletic group with high posterior probabilities (0.99) in our Bayesian analysis, as well as the parsimony analysis under equal weights. These two analyses recovered a monophyletic Lagonomegopidae, but neither of them resolved its phylogenetic placement.

The consensus trees obtained in the implied weights analysis with K = 3.505, 4.075, 4.793, 5.727, 6.991, 8.795, 11.585, which shared identical topological structures, were the ones with the highest average SPR similarity (Table 1). Analysis under a K value of 3.505 resulted in ten most fit trees with 214 steps (CI = 41, RI = 62), the strict consensus tree was used as our working hypothesis tree (Fig. 1). Symmetric resampling and Bremer support values of each nodes, as well as the sensitivity (*sensu* Giribet, 2003) to weighting regimes, are shown in this tree. The preferred tree with characters marked for exploring synapomorphies can be found in the Supporting

	K	Average SPR similarity
k1	1.432	0.811
k2	1.615	0.811
k3	1.822	0.811
k4	2.060	0.811
k5	2.336	0.897
k6	2.659	0.897
k7	3.043	0.897
k8	3.505	0.915
k9	4.075	0.915
k10	4.793	0.915
k11	5.727	0.915
k12	6.991	0.915
k13	8.795	0.915
k14	11.585	0.915
k15	16.466	0.887
k16	27.205	0.887

**Table 1.** Average similarity between trees calculated through SPR distances. The trees with the highest average similarity with the remaining trees are in **bold** 

Information (Fig. S3). Although the relationships within Lagonomegopidae are not resolved, this analysis also recovered Lagonomegopidae as a monophyletic group. A monophyletic extant Palpimanoidea is recovered here as the sister-group of Lagonomegopidae. But this clade, Lagonomegopidae + extant Palpimanoidea, has low supports (symmetric resampling = 20; Bremer support = 0.02) and only shares three homoplastic synapomorphies (characters 19, 20, 58). Entelegynae is also recovered as a monophyletic group but with low support (symmetric resampling = 22; Bremer support = 0.02). Entelegynae and the clade of Lagonomegopidae + extant Palpimanoidea are recovered as sister-groups, sharing three synapomorphies (characters 7, 47, 76). In addition, this analysis recovered both Araneoidea and the RTA clade as monophyletic but with low support.

#### SYSTEMATIC PALAEONTOLOGY

#### **ORDER ARANEAE CLERCK**, 1757

# FAMILY LAGONOMEGOPIDAE ESKOV & WUNDERLICH, 1995

GENUS SCOPOMEGOPS GUO ET AL. gen. nov.

Zoobank registration: urn:lsid:zoobank. org:act:A8809BE3-D927-432B-9C04-0AC83C27391F

*Etymology:* The genus name is the combination of *scopa*, broom in Latin, after the clustered clavate spicules on the palpal retrolateral tibial apophysis (RTA), and the Greek suffix - $\mu\epsilon\gamma\delta\psi$ , magnification, from *Lagonomegops*, type genus of the family. The name is masculine.

#### Type species: Scopomegops fax sp. nov.

*Diagnosis:* Male palpal tibia with a small apophysis on the distal retrolateral surface, several clustered clavate spicules situated on the top of retrolateral apophysis; leg metatarsus distinctly much longer than tarsus.

Remarks: CNU-ARA-MA2020001 is an adult male. It can be easily distinguished from other male lagonomegopids by the apophysis on the distal retrolateral surface of the male palpal tibia. It cannot be included in the two lagonomegopid genera (Lagonoburmops Wunderlich, 2012 and Picturmegops Wunderlich, 2015) comprised only of female specimens, because of: carapace piriform in outline, much longer than wide (carapace distinctly narrowed posteriorly, slightly wider than long in *Picturmegops*); leg setae not long and dense (legs densely covered with long setae in Lagonoburmops). Conspecific spiders have some morphological differences at different developmental stages (Foelix, **2011**). It is difficult to judge whether an adult male and a juvenile are the same species, especially in fossils. Therefore, we have erected a new genus Scopomegops based on CNU-ARA-MA2020001.

Besides Scopomegops, seven species belonging to five genera of Lagonomegopidae are known to have one or more apophyses on the male palpal tibia. In Albiburmops annulipes Wunderlich, 2017, the carapace length is equal to its width, whereas the carapace is distinctly much longer than wide in Scopomegops. In Archaelagonops propinguus Wunderlich, 2015 and Archaelagonops scorsum Wunderlich, 2015, the palpal tibia bears two apophyses, whereas there is only one in Scopomegops. In Cymbiolagonops cymbiocalcar Wunderlich, 2015, the palpal tibia bears a divided apophysis, whereas the RTA is undivided in Scopomegops. In ?Parviburmops bigibber Wunderlich, 2017, the tibial apophysis is situated on the ventral surface, but is on the retrolateral surface in Scopomegops. In *Paxillomegops longipes* Wunderlich, 2015 and ?Paxillomegops brevipes Wunderlich, 2015, the palpal tibia bears numerous short peg bristles, which are lacking in Scopomegops.

# SCOPOMEGOPS FAX GUO ET AL. sp. nov. (FIG. 2)

Zoobank registration: urn:lsid:zoobank. org:act:A897B322-B55F-4652-AF1C-FAD42209F78C

*Etymology:* The specific name is the Latin noun *fax*, meaning torch, referring to the shape of the male palp.

Holotype: Male, specimen no. CNU-ARA-MA2020001.



**Figure 1.** Strict consensus of ten trees obtained under implied weights analysis with K values of 3.505 (topology identical to K values of 4.075, 4.793, 5.727, 6.991, 8.795 and 11.585). Symmetric resampling values are shown above branches, Bremer supports below branches. Sensitivity to weighting regimes: black squares are recovered topologies, white squares are unrecovered topologies.

*Locality and horizon:* Hukawng Valley, Kachin State, northern Myanmar; lowermost Cenomanian, Mid-Cretaceous.

#### Diagnosis: As for the genus.

Taphonomic features: CNU-ARA-MA2020001: femur, patella, tibia, metatarsus, tarsus of left leg II and metatarsus, tarsus of right leg I are missing; left leg I, caused by developmental malformation or autotomy, is much smaller than right leg I; the abdomen is compressed; there is a large bubble on the left side of the abdomen, and several small bubbles around the spider. A thrips near the left leg IV is present as a syninclusion.

Description: Body length 4.97, carapace light-coloured medially, dark-coloured laterally, a pair of indistinct dark longitudinal stripes on abdomen dorsally. Carapace piriform in outline, length 2.44, width 1.54 at widest point, covered with short, dense, feathery setae pressed flat against the cuticle. Cephalic region of carapace slightly raised, with a pair of distinct protrusions beneath the inner side of PME (Fig. 2G). Fovea oval, shallow, situated in the centre of carapace, next to the posterior margin of cephalic region. Eight eyes present; PME enormous, 0.43 in diameter, situated on anterolateral corner of carapace; PLE small, placed at the mediolateral margin of cephalic region, separated 0.50 from the PME centre; AME and ALE contiguous, situated between the PME and the clypeal margin (Fig. 2G). Chelicera length 0.96, and width 0.36 at base, cheliceral insertion close to mouthparts, stridulatory files absent. Fang length 0.33. Chelicera with about six peg teeth on promargin (Fig. 2C); retromargin unrecognizable. Labium ligulate, longer than wide, not fused to sternum. Endites elongated, converging and meeting in midline; apex and prolateral margin of endites brushy with dense setae; serrula as a single row of teeth. Sternum shield shaped, covered with setae, slightly convex and without tubercles.

Palpal podomere lengths: fe 1.01, pa 0.40, ti 0.43, ta 0.98. Palp hairy, feathery setae at least present on tibia. Tibia with at least three dorsal trichobothria in a single row; a small apophysis present on the distal retrolateral surface of tibia, several clustered clavate spicules situated on the top of retrolateral apophysis (Fig. 2D, H). Male palp torch-like, cymbium elongate; embolus curved, slightly embraced by the long sheet conductor; median apophysis horn-like; subtegulum round, with a distally curved apophysis (Fig. 2I–L; File S4).

Legs long but no legs enlarged; metatarsus distinctly much longer than tarsus in all legs. Leg formula II > I > IV > III: leg I cx 0.64, tr 0.11, fe 2.81, pa 0.92, ti > 2.69; leg II cx 0.60, tr 0.11, fe 2.95, pa 0.84, ti 3.42, mt 2.42, ta 0.87; leg III cx 0.60, tr

0.10, fe 1.94, pa 0.65, ti 1.92, mt 1.13, ta 0.63; leg IV cx 0.65, tr 0.12, fe 2.51, pa 0.77, ti 2.28, mt 2.01, ta 0.82. Metatarsus and tarsus without scopulae. Distal preening comb composed of eight short macrosetae, present on metatarsus of posterior legs ventrally (Fig. 2E, F). Feathery setae at least present on all tibia. Tibiae with about 14, metatarsi with about ten, tarsi with about ten trichobothria (Fig. 2E, F). Three tarsal claws, paired claws with about four to five teeth, distal two teeth distinctly larger than others, median claw hook-like. Abdomen length 2.17, width > 0.66, densely covered with short setae. Four spinnerets visible, details unrecognizable.

*Remarks:* Holotype with left leg I much smaller than right leg I. It was caused by developmental malformation or autotomy. Autotomy is a voluntary act allowing most spiders to amputate one of their own legs to escape from a perilous situation. Although the lost legs can be replaced by new ones regenerated after the next moult, these are usually smaller, and need further moults to grow into normal size (Foelix, 2011).

# GENUS *HIATOMEGOPS* GUO *ET AL.* gen. nov.

Zoobank registration: urn:lsid:zoobank. org:act:F0561F03-E69D-45E6-A7FB-4A4D3BFEC6EE

*Etymology:* The genus name is the combination of *hiatus*, opening in Latin, after the broad tracheal spiracle behind the epigastric furrow, and *Lagonomegops*, type genus of the family, from Greek - $\mu\epsilon\gamma\delta\psi$ , magnification. The name is masculine.

#### Type species: Hiatomegops spinalis sp. nov.

*Diagnosis:* Palpal tibia and tarsus with several dorsal trichobothria; leg metatarsus slightly longer than tarsus; broad tracheal spiracle situated on postgastric area behind epigastric furrow.

*Remarks:* Finding a fossil spider species in which both males and females have been described is rare. Among 19 reported genera in Lagonomegopidae, only *Archaelagonops* Wunderlich, 2012 is known from both male and female specimens. The two female specimens, which were subsequently included in *Archaelagonops* (erected on a male specimen originally), lack sufficient evidence to belong in that genus (Wunderlich, 2012, 2015). In general, female spiders are a larger size than conspecific males. Four lagonomegopid species, *Lineaburmops beigeli, Lineaburmops maculatus* Wunderlich, 2015, *Parviburmops brevipalpus* Wunderlich, 2015 and *Planimegops parvus* Wunderlich, 2017, comprised only



**Figure 2.** *Scopomegops fax*, holotype CNU-ARA-MA2020001: A, habitus, lateral view; B, schematic drawing of lateral habitus; C, distal part of chelicerae, dorsal view, showing peg teeth; D, tibia and male palp of right pedipalp, dorsal view, showing retrolateral tibial apophysis (arrow); E, tarsus of left leg IV, lateral view, showing trichobothria (asterisks); F, metatarsus of left leg IV, lateral view, showing trichobothria (asterisks) and preening comb (arrow); G, cephalothorax, lateral view, showing eyes and abnormal left leg I; H, tibia of left pedipalp, lateral view, showing trichobothria (asterisks) and retrolateral tibial apophysis (arrow); I, left male palp, retrolateral view; J, left male palp, ventral view; K, 3D CT reconstruction of left male palp, ventral view; L, reconstruction of left male palp, ventral view. Scale bars represent 1 mm (A, B) 0.5 mm (G), 0.2 mm (D, E, F) and 0.1 mm (C, H, I, J, K, L).

of male specimens, were described as having a similar or smaller size with CNU-ARA-MA2020002, an adult female. CNU-ARA-MA2020002 can be distinguished from them by: carapace without white patch (carapace with distinct white patches in *Lineaburmops beigeli* and *Lineaburmops maculatus*); carapace much longer than wide (carapace length equal with width in Parviburmops brevipalpus); cephalic region of carapace with a pair of anterolateral protrusions next to the inner side of PME (protrusions absent in *Planimegops parvus*). We erected a new genus *Hiatomegops* based on CNU-ARA-MA2020002. Hiatomegops can be distinguished from the other two lagonomegopid genera (Lagonoburmops and Picturmegops) comprised only of female specimens by: body length 2.89 (body length about 8.00 in Lagonoburmops); carapace piriform in outline, much longer than wide (carapace distinctly narrowed posteriorly, slightly wider than long in *Picturmegops*); carapace without a stripe (carapace with distinct stripes in *Picturmegops*); metatarsus of leg I slightly longer than tarsus (metatarsus of leg I about two times longer than tarsus in Lagonoburmops).

In addition, the holotypes of Grandoculus chemahawinensis Penney, 2004, ?Lagonomegops cor Pérez-de la Fuente et al., 2013, Soplaogonomegops unzuei Pérez-de la Fuente et al., 2013, Spinomegops aragonensis Pérez-de la Fuente et al., 2013 and Spinomegops arcanus Pérez-de la Fuente et al., 2013 were identified as 'juvenile or adult female' (Penney, 2004; Pérez-de la Fuente et al., 2013). Hiatomegops can be easily separated from these by: carapace piriform in outline, without neck (carapace heart-shaped when viewed from above in ?Lagonomegops cor; cephalic region constricted in a neck, with narrowest point medially in Soplaogonomegops unzuei), fovea small oval, deeply depressed (fovea absent in Spinomegops aragonensis and Spinomegops arcanus), leg I not enlarged, metatarsus and tarsus without scopulae (leg I distinctly longer and more robust than the others, metatarsus and tarsus with scopulae hairs in Grandoculus chemahawinensis).

# HIATOMEGOPS SPINALIS GUO ET AL. sp. nov.

# (FIG. 3)

Zoobank registration: urn:lsid:zoobank. org:act:9168FFD9-29C1-4158-878B-D29E71594D41

*Etymology:* The specific name is the Latin adjective *spinalis*, meaning with spine(s), referring to the long macrosetae on the distal prolateral surface of palpal tibia.

Holotype: Female, specimen no. CNU-ARA-MA2020002.

Locality and horizon: Hukawng Valley, Kachin State, northern Myanmar; lowermost Cenomanian, Mid-Cretaceous.

#### Diagnosis: As for the genus.

Taphonomic features: CNU-ARA-MA2020002: completely preserved; many bubbles and impurities are present around the spider. Parts of insect legs are present as syninclusions.

Description: Body length 2.89, carapace and legs darkcoloured, abdomen light-coloured, with an indistinct median dorsal stripe (Fig. 3A, B). Carapace piriform in outline, length 1.30, width 0.87 at widest point, covered with short, dense, feathery setae pressed flat against the cuticle (Fig. 3A). Cephalic region of carapace slightly raised, with a pair of anterolateral protrusions next to the inner side of PME (Fig. 3E). Fovea small oval, deeply depressed, situated in the centre of carapace, next to the posterior margin of cephalic region (Fig. 3A). Eight eyes present; PME enormous, 0.26 in diameter, situated on anterolateral corner of carapace; PLE small, placed at the mediolateral margin of cephalic region, separated 0.31 from the PME centre; AME and ALE contiguous, situated between the PME and the clypeal margin. Chelicera length 0.62, width 0.27 at base, cheliceral insertion close to mouthparts, stridulatory files absent. Fang length 0.22. Chelicera with about six peg teeth on promargin (Fig. 3E), retromargin not visible. Labium ligulate, slightly longer than wide, not fused to sternum. Endites elongated, converging and meeting in midline; serrula as a single row of teeth. Sternum shield shaped, covered with setae, convex and without tubercles.

Palpal podomere lengths: fe 0.34, pa 0.17, ti 0.24, ta 0.45. Palp hairy; tibia and tarsus with several dorsal trichobothria; tibia with a long macrosetae on the distal prolateral surface (Fig. 3F). Palpal claw absent.

Legs hairy, no legs enlarged; metatarsus slightly longer than tarsus. Leg formula II > I > IV > III: leg I cx 0.23, tr 0.07, fe 0.90, pa 0.47, ti 0.76, mt 0.64, ta 0.49; leg II cx 0.26, tr 0.07, fe 0.98, pa 0.40, ti 0.93, mt 0.63, ta 0.46; leg III cx 0.27, tr 0.07, fe 0.80, pa 0.36, ti 0.58, mt 0.47, ta 0.42; leg IV cx 0.33, tr 0.09, fe 0.93, pa 0.33, ti 0.73, mt 0.67, ta 0.47. Metatarsus and tarsus without scopulae. Distal preening comb composed of seven to ten short macrosetae, present on metatarsus of posterior legs ventrally. Feathery setae at least present on femur and patella of leg II. Tibiae with about ten, metatarsi with about six, tarsi with about six trichobothria in at least two rows. Three tarsal claws, paired claws with three to five teeth, median claw hook-like.

Abdomen ovoid, length 1.33, width 0.89, densely covered with short, feathery setae. Female genitalia



**Figure 3.** *Hiatomegops spinalis*, holotype CNU-ARA-MA2020002: A, habitus, dorsal view, showing fovea (arrow) and feathery setae (magnified at the top right) on carapace; B, schematic drawing of dorsal habitus; C, habitus, ventral view; D, schematic drawing of ventral habitus; E, cephalothorax, anterior view, showing peg teeth (magnified at the bottom left); F, tibia and tarsus of right pedipalp, lateral view, showing trichobothria (asterisks) and long macrosetae (arrow); G, basal part of abdomen, ventral view, showing female genitalia and tracheal spiracle; H, schematic drawing of basal ventral abdomen; I, spinnerets, ventral view; J, schematic drawing of spinnerets, showing spigots (lower arrow). Scale bars represent 0.5 mm (A, B, C, D), 0.2 mm (E) and 0.1 mm (F, G, H, I, J).

simple, epigyne absent. A broad subovoid opening, interpreted as tracheal spiracle, situated on the postgastric area behind the epigastric furrow (Fig. 3G, H). Six spinnerets; ALS with three segments, basal segment much longer than others; PLS about as large as ALS, with two segments, distal segment almost as the equal length with basal segment; retrolateral surface of distal segment with a series of large spigots (Fig. 3I, J).

*Remarks:* Lagonomegopidae was once thought to have a female-biased sex ratio (Pérez-de la Fuente *et al.*, 2013), but as more and more male lagonomegopids are found and described, this issue was resolved (Park *et al.*, 2019). The well-preserved female genitalia of CNU-ARA-MA2020002 shows that it is an unequivocal adult female.

GENUS *LINEABURMOPS* WUNDERLICH, 2015 *Type species: Lineaburmops beigeli* Wunderlich, 2015.

# LINEABURMOPS BEIGELI WUNDERLICH, 2015

# (FIGS 4, 5)

Wunderlich (2015): 255-256, figs 236, 237, photos 104, 105.

*Material:* Two adult males, specimen nos CNU-ARA-MA2020003 and CNU-ARA-MA2020004.

*Locality and horizon:* Hukawng Valley, Kachin State, northern Myanmar; lowermost Cenomanian, Mid-Cretaceous.

Taphonomic features: CNU-ARA-MA2020003: the pedicel and abdomen are somewhat broken; metatarsus, tarsus of left leg I, II and patella of left leg IV are missing; there are a few bubbles and impurities around the spider. A thrips near the tibia of left leg III is present as a syninclusion. CNU-ARA-MA2020004: the tarsus of left leg IV is incomplete, tarsal claws are missing; and there are many impurities and cracks around the spider.

*Description:* Body length 2.98–3.08, dark-coloured, with a subtriangular white patch on clypeus, a pair of reniform white patches on the thoracic region and a pair of white longitudinal stripes on the abdomen dorsally (Figs 4A, B, 5F). Carapace length 1.57–1.59, width 0.96–0.99 at widest point, covered with short, dense, feathery setae pressed flat against the cuticle. Cephalic region of carapace slightly raised, with a pair of anterolateral protrusions next to the inner side of PME. Neck present between cephalic and thoracic regions, slightly narrower than cephalic region.

Fovea absent. Eight eyes present; PME enormous, 0.17 in diameter, situated on anterolateral corner of carapace; PLE small, placed at the mediolateral margin of cephalic region, separated 0.25 from the PME centre; AME and ALE contiguous, situated between the PME and the clypeal margin. Chelicera length 0.58 and width 0.29 at base, cheliceral insertion distinctly separated from mouthparts, stridulatory files absent. Fang length 0.19. Chelicera with about eight peg teeth on promargin (Fig. 4C); retromargin not visible. Labium subtriangular, slightly longer than wide, not fused to sternum. Endites elongated, converging but not meeting in midline; serrula as a single row of teeth. Sternum covered with setae, convex and without tubercles.

Palpal podomere lengths: fe 0.51, pa 0.18, ti 0.27, ta 0.67. Palp hairy, feathery setae at least present on tarsus. Tibia with three dorsal trichobothria in a single row (Fig. 5A). Cymbium elongate, finger-like distally (Fig. 5A). Detailed structures of male palp, such as embolus and conductor, not recognizable.

Legs long but no legs enlarged; metatarsus distinctly much longer than tarsus in anterior legs; metatarsus slightly longer than tarsus in posterior legs. Leg formula I > II > IV > III: leg I cx 0.31, tr 0.07, fe 1.47–1.58, pa 0.34, ti 1.44–1.60, mt 1.10–1.16, ta 0.66–0.75; leg II cx 0.37, tr 0.07, fe 1.49–1.53, pa 0.31-0.37, ti 1.43-1.45, mt 1.08-1.11, ta 0.66-0.71; leg III cx 0.24–0.28, tr 0.06, fe 1.14–1.17, pa 0.30–0.32, ti 1.02-1.14, mt 0.67-0.69, ta 0.50-0.57; leg IV cx 0.29-0.31, tr 0.08, fe 1.46–1.51, pa 0.32–0.35, ti 1.28–1.42, mt 0.83-0.86, ta 0.64-0.67. Metatarsus and tarsus without scopulae. Distal preening comb composed of six to eight short macrosetae, present on metatarsus of posterior legs ventrally (Fig. 5C). Feathery setae at least present on femur, patella, tibia and metatarsus (Fig. 5B). Tibiae at least with five, metatarsi at least with six, tarsi at least with nine trichobothria (Fig. 5C, D). Three tarsal claws, paired claws with five to eight teeth, median claw hook-like. Serrate accessory claw setae near the median claw present at least on left leg III (Fig. 5E).

Abdomen ovoid, length 1.27–1.34, width 0.65– 0.68, densely covered with short feathery setae. Spinnerets and anal tubercle encircled by a quadrate glabrous band. Four spinnerets visible. ALS with two segments, distal segment much smaller than basal segment; PLS about as large as ALS, details not visible (Fig. 4E, F).

*Remarks:* These two specimens can be assigned to *Lineaburmops beigeli*, the type species of genus *Lineaburmops*, by the following characters: the shape and placement of white patches on carapace and abdomen; the length of the podomeres.



**Figure 4.** *Lineaburmops beigeli*; CNU-ARA-MA2020003: A, habitus, dorsal view; B, schematic drawing of dorsal habitus; C, habitus, ventral view, showing peg teeth (magnified at the top); D, schematic drawing of ventral habitus; E, distal part of abdomen, ventral view, showing spinnerets and anal tubercle; F, schematic drawing of distal ventral abdomen. Scale bars represent 0.5 mm (A, B, C, D) and 0.1 mm (E, F).



**Figure 5.** *Lineaburmops beigeli*; CNU-ARA-MA2020003: A, tibia and male palp of right pedipalp, dorsal view, showing trichobothria (asterisks); B, distal femur of left leg III, lateral view, showing feathery setae; C, metatarsus of right leg IV, lateral view, showing trichobothria (asterisks) and preening comb (arrow); D, tarsus of left leg III, lateral view, showing trichobothria (asterisks); E, distal tarsus of left leg III, lateral view, showing claws and serrate accessory claw setae (arrow). CNU-ARA-MA2020004: F, overall habitus, dorsal view. Scale bars represent 0.5 mm (F), 0.1 mm (A, B, C, D) and 0.05 mm (E).

GENUS ODONTOMEGOPS GUO & SELDEN, 2020 Type species: Odontomegops titan Guo & Selden, 2020.

# ODONTOMEGOPS TITAN GUO & SELDEN, 2020 (FIG. 6)

Guo et al. (2020): 2–4, figs 1, 2. Material: Male, specimen no. CNU-ARA-MA2019001 (holotype). Locality and horizon: Hukawng Valley, Kachin State, northern Myanmar; lowermost Cenomanian, Mid-Cretaceous.

*Description:* See Guo *et al.* (2020). Epigastric plate with more than 30 epiandrous spigots scattered along median margin of epigastric furrow (Fig. 6C). Tracheal spiracle broad, situated on postgastric area behind epigastric furrow, immediately dividing internally into two tracheae (Fig. 6A, B).



**Figure 6.** *Odontomegops titan*; holotype CNU-ARA-MA2019001: A, basal part of abdomen, ventral view, showing male genitalia and tracheal spiracle; B, schematic drawing of basal ventral abdomen; C, male genitalia, ventral view, showing epigastric furrow and epiandrous spigots (arrows). Scale bars represent 0.5 mm (A, B) and 0.1 mm (C).

*Remarks:* After regrinding and repolishing the holotype of *Odontomegops titan*, the male genitalia and tracheal spiracle are now shown clearly. They differ from the single, broad opening in *Hiatomegops spinalis* (Fig. 3G, H), in that two small openings are situated on the postgastric area behind the epigastric furrow in *Odontomegops titan* (Fig. 6A, B). We interpret the abdomen of CNU-ARA-MA2019001 as being somewhat modified and broken, thus exposing the internal part of the tracheal spiracle. Hence, we interpret the posterior respiratory system of *Odontomegops titan* as a single, broad tracheal spiracle, which immediately divides internally into two smaller tracheae.

## DISCUSSION

## PHYLOGENETIC PLACEMENT OF LAGONOMEGOPIDAE

In our analysis, a monophyletic Lagonomegopidae is recovered but with the interfamilial relationships unclear, because the characters used here are selected for resolving the relationships among higher groups instead of lagonomegopid genera or species. The main lagonomegopid synapomorphies are: PME larger than all other eyes (character 3) and positioned on the anterolateral flanks of the carapace (character 4). Large PME also appear independently in Deinopidae Koch, 1850 and Lycosidae Sundevall, 1833 in this study, but these are not positioned in the anterolateral flanks of the carapace. Moreover, some of the potential synapomorphies with homoplasy supporting this clade are the presence of tarsal trichobothria (character 52), leg metatarsus with three or more trichobothria (character 54) and the presence of feathery setae (character 57).

Consistent with the results of previous analyses (Wood *et al.*, 2012, 2013, 2018; Fernández *et al.*, 2018; Selden *et al.*, 2020), extant Palpimanoidea, including five families, is recovered as monophyletic in the present study. The synapomorphies shared by extant palpimanoid families have been discussed before (Wood *et al.*, 2012: appendix 3). Phylogenetic relationships among extant palpimanoid families are still unstable; the previous studies (Wood *et al.*, 2012, 2013, 2018; Fernández *et al.*, 2018; Selden *et al.*, 2020) and our analysis have yielded mixed results. It should be noted, especially, that only the analysis performed by Wood *et al.* (2018) was designed for examining relationships among palpimanoids, whereas other studies, including ours, focused on different issues.

Lagonomegopidae is recovered as sister-group to extant Palpimanoidea in the present study, but the supports for this topology are weak and only three homoplastic synapomorphies are shared by Lagonomegopidae and extant Palpimanoidea: true teeth situated only on the retromargin of chelicera (character 19), chelicera with peg teeth (character 20) and a reduction in leg spination (character 58). Although regarded as a homoplastic synapomorphy, character 19 is inapplicable to the members of Huttoniidae, Mecysmaucheniidae and Stenochilidae. The presence of peg teeth on the chelicera and the spineless legs were considered as synapomorphies of Palpimanoidea in a previous study (Wood *et al.*, 2012: appendix 3). These two characters were also used by Eskov & Wunderlich (1995) to place Lagonomegopidae into Palpimanoidea. Peg teeth are not only present in palpimanoids (although absent in stenochilids), but also have evolved independently in some mimetids and malkarids; spineless legs also appear in some non-palpimanoid groups. These two characters are considered homoplastic synapomorphies shared by extant Palpimanoidea and Lagonomegopidae in this analysis.

Entelegynae is also recovered as a monophyletic group but with low support. The only non-homoplastic synapomorphy is the entelegyne female genitalia. A clade including Lagonomegopidae + extant Palpimanoidea and Entelegynae is recovered to be monophyletic with Lagonomegopidae + extant Palpimanoidea as the sister-group of Entelegynae. This clade is supported by three non-homoplastic synapomorphies: indirect eves with canoe-shaped tapetum (character 7), tarsal organ capsulate (character 47) and spinnerets with cylindrical gland spigots (character 76), although these characters are unknown for most of lagonomegopid taxa in this study. Palpimanoidea was also considered as a possible sister-group of Entelegynae in the previous studies (Wood et al., 2012, 2013, 2018; Wheeler et al., 2017; Fernández et al., 2018).

To explore the phylogenetic placement of Lagonomegopidae, we sampled six taxa representing this extinct family. These six representatives can cover most, but not all, characters present in Lagonomegopidae. The presence of a scopulae on leg I is an important feature in extant palpimanoid families. In Lagonomegopidae, four species (Archaelagonops scorsum, Grandoculus chemahawinensis, Parviburmops brevipalpus and *Paxillomegops brevipes*) have been described as having this character (Penney, 2004; Pérez-de la Fuente et al., 2013; Wunderlich, 2015), but none of these were included in our study. It is important to point out that the images showing scopulae were provided only in Grandoculus chemahawinensis (Penney, 2004; Pérez-de la Fuente et al., 2013), and these hook-tipped scopulae have a different structure from the common scopulae with spatulate tips that are present in the extant palpimanoid families (Forster & Platnick, 1984). In addition, no fossil taxa, except lagonomegopids, were included in the present study. Further phylogenetic analyses with the inclusion of more fossil spiders as terminal taxa may extend our knowledge of the evolution of the palpimanoids in the future.

EXTERNAL SEXUAL ORGANS AND APOPHYSIS ON THE RETROLATERAL SURFACE OF MALE PALPAL TIBIA (RTA) There are two general types of female spider genitalia: haplogyne and entelegyne. Haplogyne female genitalia have only a single genital opening functioning as both copulatory duct and fertilization duct, while the entelegyne female genitalia have two different forms of ducts used for copulation and oviposition, respectively (Foelix, 2011). Corresponding to the different types of female genitalia, male palps also have various morphologies. Generally, spiders with haplogyne female genitalia have simple forms of male palps, while the ones with entelegyne female genitalia have more complex male palps (Foelix, 2011). The entelegyne female genitalia was generally considered as an identified character for Entelegynae (Wheeler et al., 2017), while the haplogyne female genitalia could be found in the other spider clades. However, recent studies have suggested that the conventional model of entelegyne reproduction needs redefinition (Zhan et al., 2019) and that the entelegyne condition has evolved at least six times independently within spiders (Michalik et al., 2019). Anyway, the two general female genital types have played an important role in the classification and phylogeny of spiders.

The detailed morphological study on the external sexual organs of Lagonomegopidae is useful for examining its phylogenetic placement. Reconstruction of the male palp of *Scopomegops fax*, based on 3D CT data and optical images provides more information for our phylogenetic analysis. The holotype of *Hiatomegops spinalis* is one of the rare female lagonomegopid specimens, its female genitalia, without an epigynum, shows a relatively simple structure. But the criteria for determining the type of female genitalia is its interior structure, not just the presence or absence of epigynum. Hence, we considered the type of lagonomegopid female genitalia as unknown in this study.

Some spiders have apophyses on the surface of their male palpal tibia, functioning during the mating process (Coddington & Levi, 1991; Wheeler *et al.*, 2017). These apophyses come in a variety of forms and occupy different positions, and their homologies are ambiguous and controversial (Coddington, 1990; Griswold *et al.*, 2005; Ramírez, 2014). Here, we recognize that a single tibia may have apophyses on as many as four surfaces (prolateral, dorsal, retrolateral and ventral) and, therefore, divide them into four homology hypotheses in this study, following Griswold *et al.* (2005).

Retrolateral tibial apophyses (RTAs) was usually deemed as a synapomorphy that defined the RTA clade (Wheeler *et al.*, 2017), a highly diverse group, which accounted for about 45% of the known spider species. However, the RTA also appears in some other groups, such as *Liphistius* Schiödte, 1849, *Mecysmauchenus* Forster & Platnick, 1984 and Titanoecidae Lehtinen, 1967 (Forster & Platnick, 1984; Ramírez, 2014; Xu *et al.*, 2015). In Lagonomegopidae, seven species (including *Scopomegops fax*) are known to have one or two RTAs on the male palp, which are usually small and have several clustered clavate spicules situated on the top (Wunderlich, 2015, 2017). The homology of the RTA in these different spider groups is obscure. Our analysis suggests that lagonomegopids are not closely related to the RTA clade; the similar characters, such as leg tarsal trichobothria, shared by them are possibly convergent, because they may occupy a similar niche (Penney, 2005). There is still no unambiguous fossil evidence of RTA clade spiders in the Mesozoic (Magalhaes *et al.*, 2020), although several recent studies using molecular clocks have suggested that the RTA clade had already diversified in the Mesozoic (Fernández *et al.*, 2018; Magalhaes *et al.*, 2020).

#### TRACHEAL SPIRACLE

Spiders are the only animal group that breathe simultaneously with book lungs and tracheae (Schmitz, 2016). Basal spiders (Mesothelae, Mygalomorphae and some basal Araneomorphae) have only book lungs situated on the second and third abdominal segments, while most modern spiders (Araneomorphae) have both book lungs and tracheae (Foelix, 2011; Schmitz, 2016). The tracheae have different relative size and pattern of distribution among various spider families (Schmitz, 2013), and a recent study indicated that tracheae evolved several times independently in Araneomorphae (Ramírez et al., 2021). The tracheal spiracle is the opening of the tracheae, its number and position can be used in the classification of spiders. However, the tracheal spiracle is rarely described in fossil spiders, because it is not easy to recognize. In the holotypes of Odontomegops titan and Hiatomegops spinalis, a broad opening can be observed situated on the abdomen behind the epigastric furrow. We interpreted this opening as the tracheal spiracle based on its position and presence in both male (the holotype of *Odontomegops titan*) and female (the holotype of *Hiatomegops spinalis*) spiders.

In contrast with the tracheal spiracle, broad and placed forward in Lagonomegopidae, archaeids have two tracheal spiracles, while the other four extant palpimanoid families have a single tracheae spiracle near the spinnerets. This special tracheal spiracle is also present in Anyphaenidae Bertkau, 1878, whose members are active hunters (Ramírez, 1995; Jocqué & Dippenaar-Schoeman, 2006). The similarity in the structure and position of the tracheal spiracle between Lagonomegopidae and Anyphaenidae suggests a similar function. So, the tracheal spiracle, broad and placed forward, developed tarsal trichobothria and large PME with a tapetum, indicate that lagonomegopids were active (possibly nocturnal) hunters, like anyphaenids. This is consistent with the posit of previous study based on the large PME and the short legs (Eskov & Wunderlich, 1995).

#### CONCLUSIONS

Our study is the first attempt to test the phylogenetic placement of the extinct spider family Lagonomegopidae cladistically. We reached the following conclusions: (1) Lagonomegopidae is the potential sister-group to extant Palpimanoidea; (2) although some lagonomegopids have an RTA on the male palp, our analysis suggests that they are not closely related to the RTA clade; (3) the tracheal spiracle, broad and placed forward, developed tarsal trichobothria and large PME with a tapetum, corroborate that lagonomegopids were active (possibly nocturnal) hunters. A further phylogenetic study based on a more extensive sampling (adding more fossil taxa in the analysis) may extend our knowledge of the evolution of the palpimanoids in the future.

## ACKNOWLEDGEMENTS

We thank Matthew Downen of University of Kansas, Sheng Li and Yongjie Wang of Capital Normal University for helping with the phylogenetic analysis. We are grateful to Arsalan Zolfaghari (University of Kansas) for providing help in using the micro-CT scanner at the Department of Geology in University of Kansas. We thank Fangyuan Xia for donating specimens for this study. We thank the Editorial Board of the Zoological Journal of the Linnean Society and, in particular, Dr Maarten Christenhusz and Dr Ingi Agnarsson. We express our gratitude to three anonymous reviewers for their valuable reviews of the manuscript. XG thanks Maura Selden for help during his stay in Lawrence. This research is supported by grants from the National Natural Science Foundation of China (grants 31730087, 32020103006). XG is supported by the Graduate Student Program for International Exchange and Joint Supervision at Capital Normal University (028195509100).

#### REFERENCES

Azevedo GHF, Griswold CE, Santos AJ. 2018. Systematics and evolution of ground spiders revisited (Araneae, Dionycha, Gnaphosidae). *Cladistics* **34:** 579–626.

- **Benavides LR**, **Giribet G**, **Hormiga G. 2017**. Molecular phylogenetic analysis of 'pirate spiders' (Araneae, Mimetidae) with the description of a new African genus and the first report of maternal care in the family. *Cladistics* **33**: 375–405.
- Bond JE, Garrison NL, Hamilton CA, Godwin RL, Hedin M, Agnarsson I. 2014. Phylogenomics resolves a spider backbone phylogeny and rejects a prevailing paradigm for orb web evolution. *Current Biology* 24: 1765–1771.
- Bremer K. 1994. Branch support and tree stability. *Cladistics* 10: 295–304.
- **Coddington JA. 1990.** Cladistics and spider classification: araneomorph phylogeny and the monophyly of orbweavers (Araneae: Araneomorphae; Orbiculariae). *Acta Zoologica Fennica* **190:** 75–87.
- Coddington JA, Levi HW. 1991. Systematics and evolution of spiders (Araneae). Annual Review of Ecology and Systematics 22: 565–592.
- Cruickshank RD, Ko K. 2003. Geology of an amber locality in the Hukawng Valley, northern Myanmar. *Journal of Asian Earth Sciences* 21: 441–455.
- **Dimitrov D**, **Hormiga G. 2021.** Spider diversification through space and time. *Annual Review of Entomology* **66**: 11.1–11.17.
- Dimitrov D, Lopardo L, Giribet G, Arnedo MA, Álvarez-Padilla F, Hormiga G. 2012. Tangled in a sparse spider web: single origin of orb weavers and their spinning work unravelled by denser taxonomic sampling. *Proceedings of the Royal Society B, Biological Sciences* 279: 1341–1350.
- Dimitrov D, Astrin JJ, Huber BA. 2013. Pholcid spider molecular systematics revisited, with new insights into the biogeography and the evolution of the group. *Cladistics* 29: 132–146.
- **Dimitrov D, Benavides LR, Arnedo MA, Giribet G, Griswold CE, Scharff N, Hormiga G. 2017.** Rounding up the usual suspects: a standard target-gene approach for resolving the interfamilial phylogenetic relationships of ecribellate orb-weaving spiders with a new familyrank classification (Araneae, Araneoidea). *Cladistics* **33**: 221–250.
- **Dunlop JA**, **Penney D**, **Jekel D. 2020.** A summary list of fossil spiders and their relatives. In: *World spider catalog, v.20.5*. Bern: Natural History Museum. Availaable online at http://wsc.nmbe.ch (accessed on 04.05.2021).
- **Eskov KY**, **Wunderlich J. 1995.** On the spiders from Taimyr ambers, Siberia, with the description of a new family and with general notes on the spiders from the Cretaceous resins. *Beiträge zur Araneologie* **4:** 95–107.
- Fernández R, Hormiga G, Giribet G. 2014. Phylogenomic analysis of spiders reveals nonmonophyly of orb weavers. *Current Biology* 24: 1772–1777.
- Fernández R, Kallal RJ, Dimitrov D, Ballesteros JA, Arnedo MA, Giribet G, Hormiga G. 2018. Phylogenomics, diversification dynamics, and comparative transcriptomics across the spider tree of life. *Current Biology* 28: 1–9.
- Foelix RF. 2011. Biology of spiders, 3rd edn. New York: Oxford University Press.
- Forster RR, Platnick NI. 1984. A review of the archaeid spiders and their relatives, with notes on the limits of the

superfamily Palpimanoidea (Arachnida, Araneae). *Bulletin* of the American Museum of Natural History **178:** 1–106.

- Gao TP, Shih CK, Ren D. 2021. Behaviors and interactions of insects in mid-Mesozoic ecosystems of northeastern China. *Annual Review of Entomology* 66: 337–354.
- Garrison NL, Rodriguez J, Agnarsson I, Coddington JA, Griswold CE, Hamilton CA, Hedin M, Kocot KM, Ledford JM, Bond JE. 2016. Spider phylogenomics: untangling the spider tree of life. *PeerJ* 4: e1719.
- **Giribet G. 2003.** Stability in phylogenetic formulations and its relationship to nodal support. *Systematic Biology* **52:** 554–564.
- **Goloboff PA. 1993.** Estimating character weights during tree search. *Cladistics* **9:** 83–91.
- Goloboff PA. 2008. Calculating SPR distances between trees. Cladistics 24: 591–597.
- Goloboff PA, Catalano SA. 2016. TNT v.1.5, including a full implementation of phylogenetic morphometrics. *Cladistics* **32**: 221–238.
- Goloboff PA, Farris JS, Källersjö M, Oxelman B, Ramírez MJ, Szumik CA. 2003. Improvements to resampling measures of group support. *Cladistics* 19: 324-332.
- Goloboff PA, Farris S, Nixon K. 2008a. TNT, a free program for phylogenetic analysis. *Cladistics* 24: 774–786.
- Goloboff PA, Carpenter JM, Arias JS, Esquivel DRM. 2008b. Weighting against homoplasy improves phylogenetic analysis of morphological data sets. *Cladistics* 24: 758-773.
- Griswold CE. 1990. A revision and phylogenetic analysis of the spider subfamily Phyxelidinae (Araneae, Amaurobiidae). Bulletin of the American Museum of Natural History 196: 1–206.
- Griswold CE, Ramírez MJ, Coddington JA, Platnick NI. 2005. Atlas of phylogenetic data for entelegyne spiders (Araneae: Araneomorphae: Entelegynae) with comments on their phylogeny. *Proceedings of the California Academy of Sciences* 56: 1–324.
- **Guo XB**, **Selden PA**, **Shih CK**, **Ren D. 2020.** Two new lagonomegopid spiders (Arachnida: Araneae) from the Mid-Cretaceous of northern Myanmar, with comments on the superfamilial placement of Lagonomegopidae. *Cretaceous Research* **106**: 104257.
- Hendrixson BE, Bond JE. 2007. Molecular phylogeny and biogeography of an ancient Holarctic lineage of mygalomorph spiders (Araneae Antrodiaetidae Antrodiaetus). Molecular Phylogenetics and Evolution 42: 738–755.
- Jocqué R, Dippenaar-Schoeman AS. 2006. Spider families of the world. Tervuren: Royal Museum for Central Africa.
- Lewis PO. 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. *Systematic Biology* **50**: 913–925.
- Li F, Li S. 2018. Paleocene–Eocene and Plio–Pleistocene sea-level changes as 'species pumps' in Southeast Asia: evidence from *Althepus* spiders. *Molecular Phylogenetics and Evolution* 127: 545–555.
- Lin XD, Labandeira CC, Shih C, Hotton LC, Ren D. 2019. Life habits and evolutionary biology of new two-winged

long-proboscid scorpionflies from Mid-Cretaceous Myanmar amber. Nature Communications 10: 1235.

- Maddison WP, Maddison DR. 2019. *Mesquite: a modular system for evolutionary analysis, v.3.6.* Available at: http://www.mesquiteproject.org. (accessed on 04.05.2021).
- Magalhaes ILF, Azevedo GHF, Michalik P, Ramírez MJ. 2020. The fossil record of spiders revisited: implications for calibrating trees and evidence for a major faunal turnover since the Mesozoic. *Biological Reviews* **95**: 184–217.
- Michalik P, Kallal R, Dederichs TM, Labarque FM, Hormiga G, Giribet G, Ramírez MJ. 2019. Phylogenomics and genital morphology of cave raptor spiders (Araneae, Trogloraptoridae) reveal an independent origin of a flowthrough female genital system. Journal of Zoological Systematics and Evolutionary Research 57: 737–747.
- Mirande JM. 2009. Weighted parsimony phylogeny of the family Characidae (Teleostei: Characiformes). *Cladistics* 25: 574–613.
- Murphy JA, Roberts MJ. 2015. Spider families of the world and their spinnerets. Norwich: British Arachnological Society.
- Nixon KC. 2002. WinClada, v.1.00.08. Program and documentation. New York: Cornell University Press.
- Opatova V, Hamilton CA, Hedin M, Montes de Oca L, Kárl J, Bond JE. 2020. Phylogenetic systematics and evolution of the spider infraorder Mygalomorphae using genomic scale data. *Systematic Biology* **69**: 671–707.
- Park T-YS, Nam, K-S, Selden PA. 2019. A diverse new spider (Araneae) fauna from the Jinju Formation, Cretaceous (Albian) of Korea. *Journal of Systematic Palaeontology* 17: 1271–1297.
- **Penney D. 2004.** Cretaceous Canadian amber spider and the palpimanoidean nature of lagonomegopids. *Acta Paleontologica Polonica* **49:** 579–584.
- **Penney D. 2005.** The fossil spider family Lagonomegopidae in Cretaceous ambers with descriptions of a new genus and species from Myanmar. *Journal of Arachnology* **33**: 439–444.
- Pérez-De La Fuente R, Saupe EE, Selden PA. 2013. New lagonomegopid spiders (Araneae: †Lagonomegopidae) from Early Cretaceous Spanish amber. *Journal of Systematic Palaeontology* 11: 531–553.
- **Polotow D, Carmichael A, Griswold CE. 2015.** Total evidence analysis of the phylogenetic relationships of Lycosoidea spiders (Araneae, Entelegynae). *Invertebrate Systematics* **29:** 124–163.
- Ramírez MJ. 1995. A phylogenetic analysis of the subfamilies of Anyphaenidae (Arachnida, Araneae). *Entomologica Scandinavica* 26: 361–384.
- Ramírez MJ. 2014. The morphology and phylogeny of dionychan spiders (Araneae: Araneomorphae). *Bulletin of the American Museum of Natural History* 390: 1–374.
- Ramírez MJ, Magalhaes ILF, Derkarabetian S, Ledford J, Griswold CE, Wood HM, Hedin M. 2021. Sequence capture phylogenomics of true spiders reveals convergent evolution of respiratory systems. *Systematic Biology* **70**: 14–20.
- Rix MG, Harvey MS, Roberts JD. 2008. Molecular phylogenetics of the spider family Micropholcommatidae

(Arachnida: Araneae) using nuclear rRNA genes (18S and 28S). *Molecular Phylogenetics and Evolution* **46:** 1031–1048.

- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Ronquist F, Teslenko M, Van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61: 539–542.
- Schmitz A. 2013. Tracheae in spiders: respiratory organs for special functions. In: Nentwig W, ed. *Spider ecophysiology*. New York: Springer, 29–39.
- Schmitz A. 2016. Respiration in spiders (Araneae). Journal of Comparative Physiology B 186: 403–415.
- Schütt K. 2000. The limits of the Araneoidea (Arachnida: Araneae). Australian Journal of Zoology 48: 135–153.
- Selden PA, Penney D. 2017. Imaging techniques in the study of fossil spiders. *Earth-Science Reviews* 166: 111–131.
- Selden PA, Zhang WW, Ren D. 2016. A bizarre armoured spider (Araneae: Tetrablemmidae) from Upper Cretaceous Myanmar amber. *Cretaceous Research* 66: 129–135.
- Selden PA, Huang DY, Garwood RJ. 2020. New spiders (Araneae: Palpimanoidea) from the Jurassic Yanliao Biota of China. Journal of Systematic Palaeontology 18: 137–185.
- Shao LL, Li SQ. 2018. Early Cretaceous greenhouse pumped higher taxa diversification in spiders. *Molecular Phylogenetics and Evolution* 127: 146–155.
- Shi GH, Grimaldi DA, Harlow GE, Wang J, Wang J, Yang MC, Lei WY, Li QL, Li XH. 2012. Age constraint on Burmese amber based on U-Pb dating of zircons. *Cretaceous Research* 37: 155–163.
- Wesołowska W. 1999. A revision of the spider genus Menemerus in Africa (Araneae: Salticidae). *Genus* 10: 251–353.
- Wheeler WC, Coddington JA, Crowley LM, Dimitrov D, Goloboff PA, Griswold CE, Hormiga G, Prendini L, Ramírez MJ, Sierwald P, Almeida-Silva L, Álvarez-Padilla F, Arnedo MA, Benavides Silva LR, Benjamin SP, Bond JE, Grismado CJ, Hasan E, Hedin M, Izquierdo MA, Labarque FM, Ledford J, Lopardo L, Maddison WP, Miller JA, Piacentini LN, Platnick NI, Polotow D, Silva-Dávila D, Scharff N, Szüts T, Ubick D, Vink CJ, Wood HM, Zhang JX. 2017. The spider tree of life: phylogeny of Araneae based on target-gene analyses from an extensive taxon sampling. *Cladistics* 33: 574–616.
- **Wood HM. 2017.** Integrating fossil and extant lineages: an examination of morphological space through time (Araneae: Archaeidae). *Journal of Arachnology* **45:** 20–29.
- **Wood HM**, **Griswold CE**, **Gillespie RG**. 2012. Phylogenetic placement of pelican spiders (Archaeidae, Araneae), with insight into evolution of the 'neck' and predatory behaviours of the superfamily Palpimanoidea. *Cladistics* 28: 598–626.
- Wood HM, Matzke NJ, Gillespie RG, Griswold CE. 2013. Treating fossils as terminal taxa in divergence time estimation reveals ancient vicariance patterns in the palpimanoid spiders. Systematic Biology 62: 264–284.
- Wood HM, Gillespie RG, Griswold CE, Wainwright PC. 2015. Why is Madagascar special? The extraordinarily slow

evolution of pelican spiders (Araneae, Archaeidae). *Evolution* **69:** 462–481.

- Wood HM, González VL, Lloyd M, Coddington J, Scharff N. 2018. Next-generation museum genomics: phylogenetic relationships among palpimanoid spiders using sequence capture techniques (Araneae: Palpimanoidea). *Molecular Phylogenetics and Evolution* 127: 907–918.
- World Spider Catalog, 2021. World spider catalog, v.21.5. Bern: Natural History Museum. Available online at: http:// wsc.nmbe.ch (accessed on 16.01.2021); doi: 10.24436/2.
- Wunderlich J. 2012. On the fossil spider (Araneae) fauna in Cretaceous ambers, with descriptions of new taxa from Myanmar (Burma) and Jordan, and on the relationships of the superfamily Leptonetoidea. *Beiträge zur Araneologie* 7: 157–232.
- **Wunderlich J. 2015.** On the evolution and the classification of spiders, the Mesozoic spider faunas, and descriptions of new Cretaceous taxa mainly in amber from Myanmar (Burma) (Arachnida: Araneae). *Beiträge zur Araneologie* **9:** 21–408.

- **Wunderlich J. 2017.** New and rare fossil spiders (Araneae) in mid Cretaceous amber from Myanmar (Burma), including the description of new extinct families of the suborders Mesothelae and Opisthothelae, as well as notes on the taxonomy, the evolution and the biogeography of the Mesothelae. *Beiträge zur Araneologie* **10:** 72–279.
- Xu X, Liu FX, Chen J, Ono H, Li DQ, Kuntner M.
  2015. A genus-level taxonomic review of primitively segmented spiders (Mesothelae, Liphistiidae). ZooKeys 488: 121–151.
- Zhan YJ, Jiang H, Wu QQ, Zhang HT, Bai ZS, Kuntner M, Tu LH. 2019. Comparative morphology refines the conventional model of spider reproduction. *PLoS One* 14: e0218486.
- Zhao ZP, Yin XC, Shih CK, Gao TP, Ren D. 2020. Termite colonies from Mid-Cretaceous Myanmar demonstrate their early eusocial lifestyle in damp wood. *National Science Review* 7: 381–390.

# SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

- Fig. S1. Strict consensus of 50 most parsimonious trees obtained under equal weights analysis.
- Fig. S2. Majority rule consensus of the trees from Bayesian analysis.
- Fig. S3. The preferred tree with characters mapped on each node.
- File S1. List of morphological character descriptions and character argumentations. Word file (.docx).

File S2. Matrix. Excel file (.xls).

File S3. Matrix. TNT file (.tnt).

File S4. Movie (.mp4). 3D CT reconstruction of left male palp of CNU-ARA-MA2020001.